

A3862

ENCYCLOPEDIA OF LIFE SCIENCES

January 2004

©Macmillan Reference Ltd

Ageing Genes: Gerontogenes

Introductory

Structures and Processes

Genetics and Molecular Biology

ageing#homeostasis#homeodynamics#lifespan#longevity assurance

Rattan, Suresh IS

Suresh IS Rattan

University of Aarhus, Aarhus, Denmark

The idea of gerontogenes is in line with the evolutionary explanation of the ageing process as being caused by the absence of maintenance and repair, rather than an active, evolutionarily selected process. The term 'gerontogenes', therefore, refers to an altered state of those genes that influence ageing and longevity.

Introduction

The highly complex phenomenon of ageing is now reasonably well understood in descriptive terms. A large body of data clearly shows that ageing has many facets. Individually no tissue, organ or system (with the exception of thymus) becomes functionally exhausted even in very old organisms, and it is their combined interaction and interdependence that determines the survival of the whole. From an evolutionary point of view, the essential lifespan of a species is the time required to fulfil the darwinian purpose of life; that is, successful reproduction and continuation of generations. The later period of survival, achieved generally in protected environments and characterized by progressive ageing, is an emergent phenomenon because of the inability of the maintenance and repair pathways to continue to work efficiently. Therefore the genes that influence ageing and longevity are those that have evolved in accordance with the life history of a species for assuring its essential lifespan. Such genes are known as longevity assurance genes or vitagenes.

Nature of Ageing Genes

Evolutionary theories of ageing and longevity discount the notions of an adaptive nature of an ageing and its selection as an advantageous trait for the individual. Furthermore, the diversity of the forms and variations in which age-related alterations are manifested suggest that the progression of ageing is not programmed or deterministic but is stochastic in nature. Yet, ageing appears to have a genetic component of some kind. The role of genes in ageing is indicated by: (1) an apparent limit to maximum achievable lifespan within a species; (2) some heritability of

lifespan as evident from studies on twins; (3) human genetic mutants of premature ageing syndromes; and (4) some gene association with extreme longevity.

The paradoxical situation of the genetic aspects of ageing and longevity on the one hand, and the stochastic nature of the progression of the ageing phenotype on the other, can be resolved by developing radically novel views about the nature of [genes](#) for ageing, termed gerontogenes.

However, the proposed term gerontogenes does not refer to any 'real' genes evolved specifically to cause ageing, but to an altered state of other genes, giving the appearance of being the genes for ageing. Most importantly, the notion of [gerontogenes](#) as altered [vitagenes](#) is in line with the evolutionary explanation of the ageing process as being an emergent phenomenon caused by the absence of eternal maintenance and repair, instead of being an active process selected during evolution.

Candidate Gerontogenes

Two kinds of gene action are postulated to be responsible for the emergence of the ageing phenotype. The first considers the role of late-acting mutations which are already present at the time of fertilization and birth, and show their deleterious effects after the period of growth, development and maturation. The second category of gene action is referred to as the antagonistic pleiotropy, which involves genes selected for beneficial effects during early development but which have harmful effects in postreproductive life when they escape the force of natural selection. In both cases, these genes were not selected as the real genes causing ageing, but manifest themselves only as virtual gerontogenes due to their involvement in the progression of ageing.

Although the essential lifespan of different species may differ by several orders of magnitude, there is very little or no difference in the structure, function and activity of the so-called housekeeping genes involved in basic metabolic processes. However, there is enough evidence that the maintenance and repair pathways are one of the main determinants of longevity. Such evidence comes from comparative studies performed on species with widely varying lifespans, and from experiments performed to retard ageing and to prolong the lifespan. These genetic pathways include the efficiency of deoxyribonucleic acid (DNA) repair, the fidelity of genetic information transfer, the efficiency of protein degradation, the extent of cellular responsiveness to stress, and the capacity to protect from damage induced by free radicals and oxidation. Similarly, antiageing effects of several hormones, vitamins, a dipeptide carnosine, modified amino acids, and modified nucleic acid bases such as kinetin are also mainly due to the effects of these chemicals on maintaining, stimulating or recovering the efficiency of repair pathways. Genetic selection of *Drosophila* for longer lifespan also appears to work mainly through an increase in the efficiency of maintenance mechanisms, such as antioxidation potential and stress tolerance. The identification of long-lived mutants of the nematode *Caenorhabditis elegans* is also accompanied by an increased resistance to oxidative damage, an increase in the activities of superoxide dismutase and catalase enzymes, and an increase in thermotolerance.

Several putative gerontogenes have been reported for various ageing systems, including yeast, nematodes, insects and mammals, and the list is increasing all the time. Various approaches that have been used to identify such gerontogenes are:

- Searching for single-gene mutations resulting in a decrease or loss of gene function and extended longevity.
- Searching for increased stress resistance and extended longevity phenotype, and then identifying the genes involved.
- Creating gene-manipulated cells and organisms with reduced or enhanced gene activity.
- Using interference ribonucleic acid (RNAi) to neutralize partially or completely specific gene expression and select for extended longevity.
- Selective breeding of organisms for delayed reproduction and longer survival.
- Analysing the genes involved in human premature ageing syndromes and senescence-accelerated mice.

Additionally, genetic linkage studies for longevity in mice have identified major histocompatibility complex regions, and quantitative trait loci on human chromosomes 7, 10, 11, 12, 16, 18 and 19 as putative gerontogenes. In human centenarians, certain alleles of human leucocyte antigen (HLA) locus on chromosome 6, different alleles of *APO-E* and *APO-B*, and *DD* genotype of angiotensin-converting enzyme have been linked to their long lifespan.

The diversity of the genes associated with ageing and longevity of different organisms indicates that at the molecular level there are no universal pathways affecting ageing and longevity. Whereas the genes involved in repair and maintenance pathways may be important from an evolutionary point of view, each species may also have additional 'private' gerontogenic pathways which influence its ageing phenotype. However, there are at least three pathways that correlate well with extended longevity in a wide range of organisms, including yeast, nematodes, insects, rodents and, possibly, [humans](#). These are: (1) increased resistance to oxidative and thermal stress; (2) neuroendocrine regulation through reduced signalling of insulin-like peptides and growth hormone; and (3) gene silencing through alterations in nicotinamide - adenine dinucleotide (NAD)-dependent deacetylase activity. There may be more gerontogenic pathways yet to be discovered.

Gerontogene Therapy

Gerontogene therapy directed towards the overall ageing process seems to hold little promise. Even at the level of a single cell zygote, let alone the billions of cells in an adult human being, interfering with such a complex network and improving upon what is already a 'normal' combination for that particular individual (in the absence of any obvious genetic diseases) is a mission impossible. It appears that biogerontologists are beginning to narrow down the potential gerontogenic pathways that may be amenable to manipulation for achieving a healthy old age, with or without

a significant increase in lifespan. There is evidence that the above gerontogenic pathways can be effectively modulated by life-style alterations, such as calorie restriction, exercise, and, to some extent, by nutritional and pharmacological interventions.

Further Reading

Carnes BA, Olshansky SJ and Grahn D (2003) Biological evidence for limits to the duration of life. *Biogerontology* **4**: 31–45.

Hasty P, Campisi J, Hoeijmakers J, van Steeg H and Vijg J (2003) Aging and genome maintenance: lessons from the mouse? *Science* **299**: 1355–1359.

Hekimi S and Guarente L (2003) Genetics and the specificity of the aging process. *Science* **299**: 1351–1354.

Jazwinski SM (1996) Longevity, genes, and aging. *Science* **273**: 54–59.

Johnson TE, Henderson S, Murakami S *et al.* (2002) Longevity genes in the nematode *Caenorhabditis elegans* also mediate increased resistance to stress and prevent disease. *Journal of Inherited Metabolic Disease* **25**: 197–206.

Perls T, Kunkel L and Puca A (2002) The genetics of aging. *Current Opinion in Genetics and Development* **12**: 362–369.

Rattan SIS (1995) Gerontogenes: real or virtual? *FASEB Journal* **9**: 284–286.

Rattan SIS (2001) Applying hormesis in aging research and therapy. *Human and Experimental Toxicology* **20**: 281–285.

Rattan, SIS (ed.) (2003) *Modulating Aging and Longevity*. Dordrecht: Kluwer Academic Publishers.

SAGE KE: Science of Aging Knowledge Environment (2003).

[<http://sageke.sciencemag.org>] [Maintains a comprehensive list of gerontogenes.]